

# Zmeny lipidového spektra pri endokrinných ochoreniach

## Changes in the lipid spectrum in endocrinopathies

Patrícia Páleníková, Juraj Payer

V. interná klinika LF UK a UNB, Nemocnica Ružinov, Bratislava

✉ MUDr. Patrícia Páleníková, PhD. | endokrinologia@ru.unb.sk | www.unb.sk

### Kľúčové slová/Klíčová slova

endokrinopatie  
cholesterol  
HDL-cholesterol  
LDL-cholesterol  
lipidy  
triglyceridy

### Key words

endocrinopathies  
HDL-cholesterol  
cholesterol  
LDL-cholesterol  
lipids  
triglycerides

*Doručené do redakcie/*

*Doručeno do redakce/Received*

3. 5. 2017

*Prijaté po recenzii/*

*Přijato po recenzii/Accepted*

16. 5. 2017

### Abstrakt

Lipidy sú látky rastlinného alebo živočíšneho pôvodu. Chemicky sú to estery vyšších karboxylových mastných kyselín a alkoholov, málo rozpustné alebo nerozpustné vo vode. Sú dôležitou súčasťou buniek, nachádzajú sa v bunkových membránach a nervových tkanivách. Sú rozpúšťadlom vitamínov (A, D, E, K), hormónov, liečiv a farbív. Sú najdôležitejšou energetickou rezervou. Zo 100 g lipidov sa uvoľní cca 3 760 kJ (900 kcal). Lipoproteíny sú dôležitou stavebnou zložkou bunky, pomocou nich sa uskutočňuje transport všetkých lipidov v organizme. Koncentrácia krvných tukov je závislá od mnohých vonkajších a vnútorných faktorov, ako sú genetické predpoklady, množstvo a zloženie stravy, fyzická aktivita a samozrejme mnohé akútne i chronické ochorenia. Článok je zameraný na vplyv endokrinných ochorení na lipidy. V množstve literatúry sú údaje o vplyve endokrinopatií nekonzistentné, čo je spôsobené rôznymi faktormi, ako sú rozdiely v definovaní závažnosti ochorenia, trvania ochorenia, interindividuálne rozdiely, zohľadňovanie genetických faktorov, rozdiely v enviromentálnych faktoroch, ako sú diéta, pohyb a ďalšie.

### Abstract

Lipids are substances of biological origin – plants or animals. They are chemically esters of carboxyl fatty acids and alcohols, poorly soluble or insoluble in water. They are an important part of cell membrane and nerve tissues. They are a solvent of vitamins (A, D, E, K), hormones, drugs and dyes and the most important energy reserve. About 3 760 kJ (900 kcal) is released from 100 g of lipids. Lipoproteins are an important part of the cell with the main purpose to transport lipids. The plasma lipid levels depend on many external and internal factors – such as genetic assumptions, quantity and composition of the diet, physical activity and, of course, many acute and chronic diseases. The main focus of the article is to describe the effect of endocrine diseases on plasma lipid levels. Data on the effect of endocrinopathy are not consistent due to variety of factors such as differences in the severity of the disease, the duration of the disease, interindividual differences, consideration of genetic factors, differences in environmental factors such as diet, movement, and others.

## Vplyv endokrinopatií na lipidové spektrum Hyperprolaktinémia

Mechanizmus vplyvu prolaktínu nie je úplne objasnený, predpokladá sa pôsobenie cez viacero faktorov. Prolaktín pravdepodobne priamo pôsobí na lipidový metabolizmus, a to znížením aktivity LPL (lipoproteínovej lipázy) v ľudských tukových tkanivách a v plazme [1,2].

Druhým mechanizmom pôsobenia je pokles estrogénov spôsobený účinkom prolaktínu, čo má za následok eleváciu hladín LDL-cholesterolu (low density lipoprotein cholesterol, LDL-C) a zníženie HDL-cholesterolu (high density lipoprotein cholesterol, HDL-C). Tretím možným mechanizmom je fakt, že elevácia prolaktínu je spojená s obezitou, ktorá by mohla nepriaznivo vplyvať na plazmatické lipidy

[3]. A v neposlednom rade, pri veľkých prolaktinómoch, môže byť poškodená i produkcia rastového hormónu, a tak dochádza k ovplyvneniu hladiny lipidov (tab. 1) [1].

### Deficit rastového hormónu

Dyslipidémia je často pozorovaná u dospelých s deficitom rastového hormónu (RH) [4,5].

Deficit RH vedie k obezite, ktorá je dôležitým faktorom pri rozvoji dyslipidémie. Napriek tomu však aj u pacientov s deficitom RH a kontrolovaným body mass indexom je dyslipidémia stále prítomná [4].

### Efekt liečby RH

Metaanalýza Newmana potvrdila efekt nízkej aj vysokej dávky (> 0,7 mg/d) RH na lipidový status – znížil sa celkový aj LDL-C, ale zmeny v HDL-C a triglyceridov (TG) boli nesignifikantné [6].

### Mechanizmus zmien v lipidovom spektre pri deficite RH

RH zvyšuje expresiu hepatálnych LDL-receptorov [7,8]. Pacienti liečení RH majú zvýšenú expresiu LDL-receptorov, ktorá vedie k zníženiu plazmatického LDL. Avšak u pacientov s familiárnou hypercholesterolémiou zapríčinenou nefunkčnosťou LDL-receptorov liečba RH nemá úspech. Pacienti s deficitom RH majú zvýšenú hepatálnu produkciu VLDL (very low density lipoprotein cholesterol) a redukovaný klírens VLDL, ktoré majú za následok zvýšenie TG v plazme. RH stimuluje lipolýzu a zvyšuje oxidáciu mastných kyselín (tab. 2) [9].

### Akromegália

Pacienti s akromegáliou majú často pozorovanú zvýšenú hladinu TG a zníženú hladinu HDL cholesterolu [10,11,12,19,20]. Efekt akromegálie na celkový cholesterol a LDL-C je variabilný [13,14]. Liečba akromegálie, ktorá normalizuje hodnoty RH a IGF1 (insulin growth factor) vyúsťuje do zníženia TG a zvýšenia HDL-C [10,15].

Tab. 1. Vplyv zvýšených hodnôt prolaktínu na parametre lipidového spektra

T-C	↑
LDL-C	↑
HDL-C	→
TG	→

T-C – celkový cholesterol (total cholesterol)

Tab. 2. Vplyv deficitu RH na parametre lipidového spektra

T-C	↑
LDL	↑
HDL	↓
TG	↑

### Mechanizmus pôsobenia

RH má schopnosť zvyšovať lipolýzu, ktorá ponúka mastné kyseliny na syntézu TG v pečeni, a tým zvyšuje produkciu VLDL [16,17]. Niektoré štúdie ukázali, že u pacientov s akromegáliou je aktivita LPL (lipoproteínovej lipázy) znížená [14,18] a aktivita LCAT (lecithin cholesterol acyltransferáza), CEPT (cholesteryl ester transfer protein), hepatálna lipáza a fosfolipidový transfer boli znížené (tab. 3) [19,20].

### Hypotyreóza

Približne od 30. rokov 20. storočia je známe, že znížená činnosť štítnej žľazy je spojená so zvýšením plazmatických hladín cholesterolu [21]. Lipidový profil pacienta s hypotyreózou je charakterizovaný zvýšením celkového a LDL-C, pričom hladiny LDL sú zvýšené nápadne. U pacientov so signifikantnou eleváciou LDL-C by malo byť v klinickej praxi rutinným vyšetrením vyšetrenie TSH (tyreostimulačný hormón), aby sa vylúčila hypotyreóza ako príčina hypercholesterolémie. U pacientov so sekundárnou hypotyreózou nebol pozorovaný rozdiel v hodnotách lipidového spektra [22]. Po úprave tyreoidálnej funkcie je tendencia k normalizácii hodnôt lipidov [21,23]. Efekt subklinickej hypotyreózy na lipidové spektrum je variabilný a v niektorých štúdiách boli popísané zmeny podobné ako pri plne rozvinutej hypotyreóze [24,25]. Podľa novších analýz sa však nepotvrdil benefit substitučnej liečby tyroxínom na parametre lipidového spektra pri liečbe subklinickej hypotyreózy [26]. Americká tyreologická spoločnosť a Americká spoločnosť klinických endokrinológov odporúčajú liečbu subklinickej hypotyreózy tyroxínom len v prípade, ak je TSH > 10 mIU/l [27]. Pacienti so signifikantnou hyperlipidémiou majú byť liečení zmenou životného štýlu a hypolipidemikami.

### Mechanizmus zmien

Tyreoidálne hormóny regulujú expresiu a aktivitu množstva kľúčových enzýmov a receptorov, ktoré ovplyvňujú hladiny lipidov. Primárnym mechanizmom je zníženie hladiny LDL-receptorov v pečeni. Hormóny štítnej žľazy stimulujú expresiu LDL-receptorov zvýšením SREBP 2 (sterol regulatory element binding protein) alebo priamym pôsobením na receptorový promóter [28,29] a teda pri hypotyreóze je množstvo LDL-receptorov redukované, čo vedie k zníženému klírensu cirkulujúceho LDL v plazme [21,30,31,32]. Tyreoidálne hormóny tiež stimulujú konverziu cholesterolu na žlčové kyseliny zvýšením cholesterol-7 $\alpha$ -hydroxylázy, čo je iniciálny enzým syntézy žlčových kyselín [33]. Navyše,

Tab. 3. Vplyv zvýšeného RH na parametre lipidového spektra

T-C	rôzny
LDL	rôzny
HDL	↓
TG	↑

tyroidálne hormóny tiež stimulujú expresiu transportérov, ktoré sa zúčastňujú na transporte cholesterolu z hepatocytu do žlče [34]. Hypotyreóza je asociovaná so zvýšenou črevnou absorpciou cholesterolu [34]. Ako je uvedené v tab. 4, hypotyreóza má len mierny vplyv na plazmatické hladiny TG a HDL-C. Niekoľko štúdií potvrdilo, že hormóny štítnej žľazy stimulujú LPL [35,36,37] a tiež zvyšujú CEPT alebo transfer cholesterolových esterov, hepatálnu lipázu, LCAT [35,36,37,38,39].

### Hypertyreóza

Pri hypertyreóze sú cholesterol, LDL-C a HDL-C znížené [21,40]. Úprava hypertyreózy rezultuje do normalizácie parametrov lipidového spektra.

Mechanizmy účinku sú uvedené v časti o tyroidálnych hormónoch (tab. 5).

### Hyperkorticizmus

V klinickom obraze Cushingovho syndrómu je jedným z dominujúcich znakov abdominálna obezita, v laboratórnom obraze je to dyslipidémia charakterizovaná eleváciou TG a celkového cholesterolu a LDL-C [41,42,43]. Zvýšenie celkového a LDL-C koreluje so závažnosťou Cushingovho syndrómu [41,42]. Ak je prítomný i steroidný diabetes mellitus, tento môže ďalej zhoršovať lipidový status pacienta s Cushingovým syndrómom [44]. Úprava lipidového spektra sa pozoruje po liečbe a znížení hladín kortizolu [41,45]. V longitudinálnej štúdii 25 pacientov dosiahlo signifikantné zníženie hladín LDL-C po 1 roku normalizácie kortizolu, ale tieto hladiny ostávali naďalej o niečo vyššie v porovnaní s kontrolným súborom [41,45].

### Mechanizmus zmien

Presné mechanizmy ako exces glukokortikoidov ovplyvňuje zmeny v lipidovom spektre ešte neboli objasnené. V štúdií s potkanmi sa pozorovalo, že glukokortikoidy znižujú expresiu hepatických LDL-receptorov [47]. Galman pozoroval, že

ACTH (adrenokortikotropný hormón) stimulácia nadobličiek znížila expresiu ako hepatických LDL-receptorov, tak aj SR-B1 receptorov (scavengerový receptor triedy B typ I). Predpokladá, že hormóny iné ako glukokortikoidy, ktoré sú uvoľňované aktivovanými nadobličkami, môžu mať efekt na pečenné receptory [48]. Podanie glukokortikoidov stimuluje syntézu mastných kyselín zvýšením aktivity acetylCoA karboxylázy [49,50]. Glukokortikoidy tiež stimulujú enzýmy potrebné na syntézu TG v pečeni [50,51], zvyšujú expresiu TG-lipázy a hormón senzitivnej lipázy, ktoré sú kľúčové pri rozklade TG na voľné mastné kyseliny v tukovom tkanive [52]. Glukokortikoidy zvyšujú aj lipolýzu v tukovom tkanive prostredníctvom zvýšenia hladiny cAMP (cyklický adenosínmonofosfát), ktorý stimuluje aktiváciu proteín kinázy A (PKA) vedúcej k fosforylácii hormón senzitivnej lipázy a perilipínu (tab. 6) [53].

### Testosterón a lipidový metabolizmus

Mnohé epidemiologické štúdie ukázali, že sérový testosterón (TST) priamo koreluje s hladinou HDL-C a apolipoproteínu A-I [55,56,57,58]. Navyše, nízke hladiny TST nepriamo úmerne korelujú s hladinou celkového cholesterolu, apolipoproteínu B a hladinami TG [56,57]. Takže muži s nízkou hladinou TST majú proaterogénny lipoproteínový vzor s nízkym HDL-C a vysokými hladinami TG a LDL-C a sú viac postihnutí metabolickým syndrómom (tab. 7) [58,59].

### Liečba testosterónom

Metaanalýza Whitsela ukázala, že celkový cholesterol, HDL-C a LDL-C sa znižujú po intramuskulárnom podaní testosterónu, ale hladina TG sa nemení [60]. Metaanalýza Isidora tiež poukázala na zníženie HDL-C, ale nedokázala žiadne zmeny v hladine LDL-C pri intramuskulárnom a transkutánnom podaní [61]. Recentná metaanalýza Coronu nepotvrdila zmeny v hladinách HDL-C, ale poukázala na malé zníženie v hladine celkového cholesterolu a TG [62]. Z klinického hľadiska je dôležité, že pacienti

Tab. 4. Vplyv hypotyreózy na parametre lipidového spektra

	hypotyreóza	subklinická hypotyreóza
T-C	↑	→ alebo ↑
LDL-C	↑	→ alebo ↑
HDL-C	→ alebo mierne ↑	→

Tab. 5. Vplyv hypertyreózy na parametre lipidového spektra

T-C	↓
LDL-C	↓
HDL-C	↓
TG	rôzne

Tab. 6. Vplyv hyperkorticizmu na parametre lipidového spektra

T-C	↑
LDL-C	↑
HDL-C	rôzny
TG	↑

Tab. 7. Znížené hladiny TST a vplyv na lipidové spektrum

T-C	↑
HDL-C	↓
LDL-C	↑
TG	↑

s deficitom androgénov, ktorí sú liečení substitučnou dávkou testosterónu, majú len malé alebo žiadne zmeny v hladinách plazmatických lipidov a lipoproteínov [63]. Použitie vysokých dávok androgénnych steroidov u mladých mužov za účelom zvýšenia svalovej hmoty má iný vplyv na lipidové spektrum. V štúdiu Webba, v ktorej sa podávali vysoké dávky androgénov mladým mužom, pozorovali, že hladiny HDL-C sa znížili na viac ako 50 % z východiskovej hodnoty, tiež boli zaznamenané zvýšené hodnoty LDL-C [64]. Preto ak sa u mužov-športovcov zistia nečakane nízke hodnoty HDL-C, môže byť vyslovené podozrenie na zneužitie anabolických steroidov.

### Mechanizmus vplyvu testosterónu na lipidy

Zníženie HDL-C je pripisované zvýšenej expresii SR-B1 v pečeni a zvýšeniu aktivity hepatálnej lipázy. Podanie TST zvyšuje aktivitu hepatálnej lipázy, ale má len malý efekt na LPL [65,66]. Zvýšenie SR-B1 v pečeni uľahčuje transfer cholesterolu z HDL-častíc do hepatocytov. Zvýšenie aktivity hepatálnej lipázy zvyšuje hydrolýzu TG [67].

### Efekt ženských pohlavných hormónov na lipidový metabolizmus

Lipidový profil premenopauzálnych žien je menej pro-aterogénny ako u mužov [68,69,70]. Najmä hladiny HDL-C sú zvýšené (oproti mužom cca o 10 mg/dl), zatiaľ čo LDL-C je porovnateľný s hladinou u mužov. Hladiny TG sú u premenopauzálnych žien nižšie.

Zmeny v lipidovom spektre po menopauze sú relatívne malé a výsledky v literatúre pomerne variabilné [68,69,70]. Najčastejšie sa pozoruje zvýšenie hladín LDL-C. Hodnoty HDL-C sú stabilné, len niektoré štúdie pozorovali malé zníženie. Menopauza je asociovaná so zvýšením celkového a centrálneho tuku, zníženou inzulínovou senzitivitou, ktorých vplyv na lipidový metabolizmus je známy [71].

### Liečba estrogénmi

Efekt perorálnej estrogénovej liečby na lipidy je známy niekoľko rokov. Ich podanie zvyšuje hladiny HDL-C o 5–15 % a znižuje LDL-C o 5–20 % [68, 72,73]. Transdermálne podanie má menší efekt na lipidy a lipoproteíny. Efekt na TG je minimálny, a preto u pacientiek, ktoré už majú abnormality v metabolizme TG, sa preferuje použitie transdermálnych foriem [74,75].

### Kombinovaná liečba estrogénom a progesterónom

Progesterónové prípravky majú efekt na lipidy podobný androgénom – znižujú HDL-C a TG, ale majú len malý alebo žiadny efekt na hladinu LDL-C [68,72,73]. Kombinované prípravky preto zmierňujú estrogénmi podmienené zvýšenie HDL-C a TG [73,76]. Podľa Godslandovej analýzy poradie preprátatov s narastajúcim účinkom na lipidy je nasledovné: dydrogesteron a medrogestone, progesterón, cyproteron acetát, medroxyprogesteron acetát, transder-

málny norethindron acetát, norgestrel, perorálny norethindron acetát [72].

Syndróm polycystických ovárií charakterizujú nízke hladiny HDL-C, zvýšené TG a LDL-C [77,78]. Tieto zmeny sú prítomné aj u pacientiek, ktoré nie sú obézne. Abnormality v lipidovom spektre sú multifaktoriálne podmienené – zvýšením androgénov, znížením estrogénov, obezitou, inzulínovou rezistenciou, genetickými faktormi [78,78].

### Záver

Rôzne endokrinné ochorenia ovplyvňujú metabolizmus lipidov a lipoproteínov rôzne. Nebola dokázaná jasná korelácia medzi zmenami, ktoré sú spôsobené deficitom hormónov a nadbytkom hormónov alebo ich terapiou. Poruchy lipidového spektra, ktoré sa vyskytujú pri nadbytku PRL, deficite RH, Cushingovom syndróme, mužskom hypogonadizme, syndróme polycystických ovárií a estrogénnom deficite môžu vyústiť do zvýšeného aterosklerotického rizika, a tým sa zvyšuje riziko kardiovaskulárnych komplikácií. Liečbou týchto ochorení sa tieto riziká znižujú.

### Literatúra

1. Pelkonen R, Nikkilä EA, Grahne B. Serum lipids, postheparin plasma lipase activities and glucose tolerance in patients with prolactinoma. *Clin Endocrinol (Oxf)* 1982; 16(4): 383–390.
2. Ling C, Svensson L, Odén B et al. Identification of functional prolactin (PRL) receptor gene expression: PRL inhibits lipoprotein lipase activity in human white adipose tissue. *J Clin Endocrinol Metab* 2003; 88(4): 1804–1808.
3. Pala NA, Laway BA, Misgar RA et al. Metabolic abnormalities in patients with prolactinoma: response to treatment with cabergoline. *Diabetol Metab Syndr* 2015; 7: 99. Dostupné z DOI: <<http://dx.doi.org/10.1186/s13098-015-0094-4>>.
4. Rosén T, Edén S, Larson G et al. Cardiovascular risk factors in adult patients with growth hormone deficiency. *Acta Endocrinol (Copenh)* 1993; 129(3): 195–200.
5. de Boer H, Blok GJ, Voerman HJ et al. Serum lipid levels in growth hormone-deficient men. *Metabolism* 1994; 43(2): 199–203.
6. Newman CB, Carmichael JD, Kleinberg DL. Effects of low dose versus high dose human growth hormone on body composition and lipids in adults with GH deficiency: a meta-analysis of placebo-controlled randomized trials. *Pituitary* 2015; 18(3): 297–305. Dostupné z DOI: <<http://dx.doi.org/10.1007/s11102-014-0571-z>>.
7. Parini P, Angelin B, Lobie PE et al. Growth hormone specifically stimulates the expression of low density lipoprotein receptors in human hepatoma cells. *Endocrinology* 1995; 136(9): 3767–3773.
8. Rudling M, Norstedt G, Olivecrona H et al. Importance of growth hormone for the induction of hepatic low density lipoprotein receptors. *Proc Natl Acad Sci U S A* 1992; 89(15): 6983–6987.
9. Møller N, Jørgensen JO. Effects of growth hormone on glucose, lipid, and protein metabolism in human subjects. *Endocr Rev* 2009; 30(2): 152–177. Dostupné z DOI: <<http://dx.doi.org/10.1210/er.2008-0027>>.
10. Colao A, Spinelli L, Cuocolo A et al. Cardiovascular consequences of early-onset growth hormone excess. *J Clin Endocrinol Metab* 2002; 87(7): 3097–3104.
11. Colao A, Pivonello R, Grasso LF et al. Determinants of cardiac disease in newly diagnosed patients with acromegaly: results of a 10 year survey study. *Eur J Endocrinol* 2011; 165(5): 713–721. Dostupné z DOI: <<http://dx.doi.org/10.1530/EJE-11-0408>>.
12. Boero L, Manavela M, Gómez Rosso L et al. Alterations in biomarkers of cardiovascular disease (CVD) in active acromegaly. *Clin Endocrinol (Oxf)* 2009; 70(1): 88–95. Dostupné z DOI: <<http://dx.doi.org/10.1111/j.1365-2265.2008.03323.x>>.

13. Vilar L, Naves LA, Costa SS et al. Increase of classic and nonclassic cardiovascular risk factors in patients with acromegaly. *Endocr Pract* 2007; 13(4): 363–372.
14. Takeda R, Tatami R, Ueda K et al. The incidence and pathogenesis of hyperlipidaemia in 16 consecutive acromegalic patients. *Acta Endocrinol (Copenh)* 1982; 100(3): 358–362.
15. Colao A, Marzullo P, Lombardi G. Effect of a six-month treatment with lanreotide on cardiovascular risk factors and arterial intima-media thickness in patients with acromegaly. *Eur J Endocrinol* 2002; 146(3): 303–309.
16. Christ ER, Cummings MH, Albany EE et al. Effects of growth hormone (GH) replacement therapy on very low density lipoprotein apolipoprotein B100 kinetics in patients with adult GH deficiency: a stable isotope study. *J Clin Endocrinol Metab* 1999; 84(1): 307–316.
17. Moller N, Vendelbo MH, Kampmann U et al. Growth hormone and protein metabolism. *Clin Nutr* 2009; 28(6): 597–603. Dostupné z DOI: <<http://dx.doi.org/10.1016/j.clnu.2009.08.015>>.
18. Twickler TB, Dallinga-Thie GM, Zelissen PM et al. The atherogenic plasma remnant-like particle cholesterol concentration is increased in the fasting and postprandial state in active acromegalic patients. *Clin Endocrinol (Oxf)* 2001; 55(1): 69–75.
19. Boero L, Manavela M, Meroño T et al. GH levels and insulin sensitivity are differently associated with biomarkers of cardiovascular disease in active acromegaly. *Clin Endocrinol (Oxf)* 2012; 77(4): 579–585. Dostupné z DOI: <<http://dx.doi.org/10.1111/j.1365-2265.2012.04414.x>>.
20. Beentjes JA, van Tol A, Sluiter WJ et al. Low plasma lecithin:cholesterol acyltransferase and lipid transfer protein activities in growth hormone deficient and acromegalic men: role in altered high density lipoproteins. *Atherosclerosis* 2000; 153(2): 491–498.
21. Duntas LH, Brenta G. The effect of thyroid disorders on lipid levels and metabolism. *Med Clin North Am* 2012; 96(2): 269–281. Dostupné z DOI: <<http://dx.doi.org/10.1016/j.mcna.2012.01.012>>.
22. Valdemarsson S, Hedner P, Nilsson-Ehle P. Dyslipoproteinaemia in hypothyroidism of pituitary origin: effects of L-thyroxine substitution on lipoprotein lipase, hepatic lipase, and on plasma lipoproteins. *Acta Endocrinol (Copenh)* 1983; 103(2): 192–197.
23. O'Brien T, Dinneen SF, O'Brien PC et al. Hyperlipidemia in patients with primary and secondary hypothyroidism. *Mayo Clin Proc* 1993; 68(9): 860–866.
24. Pearce EN. Update in lipid alterations in subclinical hypothyroidism. *J Clin Endocrinol Metab* 2012; 97(2): 326–333. Dostupné z DOI: <<http://dx.doi.org/10.1210/jc.2011-2532>>.
25. Wiersinga WM. Adult Hypothyroidism. In De Groot LJ, Chrousos G, Dungan K (eds). *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc. 2000–2017. Dostupné z WWW: <<https://www.ncbi.nlm.nih.gov/books/NBK285561/#tyd-hypothyroidism.toc-9-1-historical>>.
26. Rugge JB, Bougatsos C, Chou R. Screening and treatment of thyroid dysfunction: an evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2015; 162(1): 35–45. <<http://dx.doi.org/10.7326/M14-1456>>.
27. Garber JR, Cobin RH, Gharib H et al. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Endocr Pract* 2012; 18(6): 988–1028.
28. Lopez D, Abisambra Socarrás JF, Bedi M et al. Activation of the hepatic LDL receptor promoter by thyroid hormone. *Biochim Biophys Acta* 2007; 1771(9): 1216–1225.
29. Shin DJ, Osborne TF. Thyroid hormone regulation and cholesterol metabolism are connected through Sterol Regulatory Element-Binding Protein-2 (SREBP-2). *J Biol Chem* 2003; 278(36): 34114–34118.
30. Walton KW, Scott PJ, Dykes PW et al. The significance of alterations in serum lipids in thyroid dysfunction. II. Alterations of the metabolism and turnover of 131-I-low-density lipoproteins in hypothyroidism and thyrotoxicosis. *Clin Sci* 1965; 29(2): 217–238.
31. Staels B, Van Tol A, Chan L et al. Alterations in thyroid status modulate apolipoprotein, hepatic triglyceride lipase, and low density lipoprotein receptor in rats. *Endocrinology* 1990; 127(3): 1144–1152.
32. Scarabottolo L, Trezzi E, Roma P et al. Experimental hypothyroidism modulates the expression of the low density lipoprotein receptor by the liver. *Atherosclerosis* 1986; 59(3): 329–333.
33. Bonde Y, Breuer O, Lütjohann D et al. Thyroid hormone reduces PCSK9 and stimulates bile acid synthesis in humans. *J Lipid Res* 2014; 55(11): 2408–2415. Dostupné z DOI: <<http://dx.doi.org/10.1194/jlr.M051664>>.
34. Gälman C, Bonde Y, Matasconi M et al. Dramatically increased intestinal absorption of cholesterol following hypophysectomy is normalized by thyroid hormone. *Gastroenterology* 2008; 134(4): 1127–1136. Dostupné z DOI: <<http://dx.doi.org/10.1053/j.gastro.2008.01.032>>.
35. Kuusi T, Taskinen MR, Nikkilä EA. Lipoproteins, lipolytic enzymes, and hormonal status in hypothyroid women at different levels of substitution. *J Clin Endocrinol Metab* 1988; 66(1): 51–56.
36. Lam KS, Chan MK, Yeung RT. High-density lipoprotein cholesterol, hepatic lipase and lipoprotein lipase activities in thyroid dysfunction—effects of treatment. *Q J Med* 1986; 59(229): 513–521.
37. Valdemarsson S, Hedner P, Nilsson-Ehle P. Reversal of decreased hepatic lipase and lipoprotein lipase activities after treatment of hypothyroidism. *Eur J Clin Invest* 1982; 12(5): 423–428.
38. Tan KC, Shiu SW, Kung AW. Plasma cholesteryl ester transfer protein activity in hyper- and hypothyroidism. *J Clin Endocrinol Metab* 1998; 83(1): 140–143.
39. Johansson L, Rudling M, Scanlan TS et al. Selective thyroid receptor modulation by GC-1 reduces serum lipids and stimulates steps of reverse cholesterol transport in euthyroid mice. *Proc Natl Acad Sci USA* 2005; 102(29): 10297–10302.
40. Heimberg M, Olubadewo JO, Wilcox HG. Plasma lipoproteins and regulation of hepatic metabolism of fatty acids in altered thyroid states. *Endocr Rev* 1985 Fall; 6(4): 590–607.
41. Arnaldi G, Scandali VM, Trentin L et al. Pathophysiology of dyslipidemia in Cushing's syndrome. *Neuroendocrinology* 2010; 92(Suppl 1): 86–90. Dostupné z DOI: <<http://dx.doi.org/10.1159/000314213>>.
42. Mancini T, Kola B, Mantero F et al. High cardiovascular risk in patients with Cushing's syndrome according to 1999 WHO/ISH guidelines. *Clin Endocrinol (Oxf)* 2004; 61(6): 768–777.
43. Colao A, Pivonello R, Spiezia S et al. Persistence of increased cardiovascular risk in patients with Cushing's disease after five years of successful cure. *J Clin Endocrinol Metab* 1999; 84(8): 2664–2672.
44. Feingold KR, Grunfeld C. Diabetes and Dyslipidemia. In De Groot LJ, Chrousos G, Dungan K et al (eds). *Endotext* [Internet]. MDText.com, Inc: South Dartmouth (MA) 2015. Dostupné z WWW: <<https://www.ncbi.nlm.nih.gov/books/NBK305900/>>.
45. Taskinen MR, Nikkilä EA, Pelkonen R et al. Plasma lipoproteins, lipolytic enzymes, and very low density lipoprotein triglyceride turnover in Cushing's syndrome. *J Clin Endocrinol Metab* 1983; 57(3): 619–626.
46. Faggiano A, Pivonello R, Spiezia S et al. Cardiovascular risk factors and common carotid artery caliber and stiffness in patients with Cushing's disease during active disease and 1 year after disease remission. *J Clin Endocrinol Metab* 2003; 88(6): 2527–2533.
47. Hazra A, Pyszczyński NA, DuBois DC et al. Modeling of corticosteroid effects on hepatic low-density lipoprotein receptors and plasma lipid dynamics in rats. *Pharm Res* 2008; 25(4): 769–780. Dostupné z DOI: <<http://dx.doi.org/10.1007/s11095-007-9371-8>>.
48. Gälman C, Angelin B, Rudling M. Prolonged stimulation of the adrenals by corticotropin suppresses hepatic low-density lipoprotein and high-density lipoprotein receptors and increases plasma cholesterol. *Endocrinology* 2002; 143(5): 1809–1816.
49. Amatruda JM, Danahy SA, Chang CL. The effects of glucocorticoids on insulin-stimulated lipogenesis in primary cultures of rat hepatocytes. *Biochem J* 1983; 212(1): 135–141.
50. Lau DC, Roncari DA. Effects of glucocorticoid hormones on lipid-synthetic enzymes from different adipose tissue regions and from liver. *Can J Biochem Cell Biol* 1983; 61(12): 1245–1250.
51. Dolinsky VW, Douglas DN, Lehner R et al. Regulation of the enzymes of hepatic microsomal triacylglycerol lipolysis and re-esterification by the glucocorticoid dexamethasone. *Biochem J* 2004; 378(Pt 3): 967–974.

52. Campbell JE, Peckett AJ, D'souza AM et al. Adipogenic and lipolytic effects of chronic glucocorticoid exposure. *Am J Physiol Cell Physiol* 2011; 300(1): C198–C209. Dostupné z DOI: <<http://dx.doi.org/10.1152/ajpcell.00045.2010>>.
53. Xu C, He J, Jiang H et al. Direct effect of glucocorticoids on lipolysis in adipocytes. *Mol Endocrinol* 2009; 23(8): 1161–1170. Dostupné z DOI: <<http://dx.doi.org/10.1210/me.2008-0464>>.
54. Peckett AJ, Wright DC, Riddell MC. The effects of glucocorticoids on adipose tissue lipid metabolism. *Metabolism* 2011; 60(11): 1500–1510. Dostupné z DOI: <<http://dx.doi.org/10.1016/j.metabol.2011.06.012>>.
55. Agledahl I, Skjaerpe PA, Hansen JB et al. Low serum testosterone in men is inversely associated with non-fasting serum triglycerides: the Tromso study. *Nutr Metab Cardiovasc Dis* 2008; 18(4): 256–262.
56. Haffner SM, Mykkänen L, Valdez RA et al. Relationship of sex hormones to lipids and lipoproteins in nondiabetic men. *J Clin Endocrinol Metab* 1993; 77(6):1610–1615.
57. Hämäläinen E, Adlercreutz H, Ehnholm C et al. Relationships of serum lipoproteins and apoproteins to sex hormones and to the binding capacity of sex hormone binding globulin in healthy Finnish men. *Metabolism* 1986; 35(6): 535–541.
58. Wu FC, von Eckardstein A. Androgens and coronary artery disease. *Endocr Rev* 2003; 24(2): 183–217.
59. Corona G, Monami M, Rastrelli G et al. Testosterone and metabolic syndrome: a meta-analysis study. *J Sex Med* 2011; 8(1): 272–83. Dostupné z DOI: <<http://dx.doi.org/10.1111/j.1743-6109.2010.01991.x>>.
60. Whitsel EA, Boyko EJ, Matsumoto AM et al. Intramuscular testosterone esters and plasma lipids in hypogonadal men: a meta-analysis. *Am J Med* 2001; 111(4): 261–269.
61. Isidori AM, Giannetta E, Greco EA et al. Effects of testosterone on body composition, bone metabolism and serum lipid profile in middle-aged men: a meta-analysis. *Clin Endocrinol (Oxf)* 2005; 63(3): 280–293.
62. Corona G, Giagulli VA, Maseroli E et al. THERAPY OF ENDOCRINE DISEASE: Testosterone supplementation and body composition: results from a meta-analysis study. *Eur J Endocrinol* 2016; 174(3): R99–R116. Dostupné z DOI: <<http://dx.doi.org/10.1530/EJE-15-0262>>.
63. MD KF, Brinton EA, Grunfeld C. The Effect of Endocrine Disorders on Lipids and Lipoproteins. In: De Groot LJ, Chrousos G, Dungan K et al (eds). *Endotext* [Internet]. MDText.com: South Dartmouth (MA) 2017. Dostupné z WWW: <<https://www.ncbi.nlm.nih.gov/books/NBK409608/#/endo-disorder-effects.toc-abstract>>.
64. Webb OL, Laskarzewski PM, Glueck CJ. Severe depression of high-density lipoprotein cholesterol levels in weight lifters and body builders by self-administered exogenous testosterone and anabolic-androgenic steroids. *Metabolism* 1984; 33(11): 971–975.
65. Friedl KE, Hannan CJ Jr, Jones RE et al. High-density lipoprotein cholesterol is not decreased if an aromatizable androgen is administered. *Metabolism* 1990; 39(1): 69–74.
66. Tan KC1, Shiu SW, Kung AW. Alterations in hepatic lipase and lipoprotein subfractions with transdermal testosterone replacement therapy. *Clin Endocrinol (Oxf)* 1999; 51(6): 765–769.
67. Feingold KR, Grunfeld C. Introduction to Lipids and Lipoproteins. In: De Groot LJ, Chrousos G, Dungan K et al (eds). *Endotext* [Internet]. MDText.com: South Dartmouth (MA) 2015. Dostupné z WWW: <[https://www.ncbi.nlm.nih.gov/books/NBK305896/#lipid\\_intro.toc-abstract](https://www.ncbi.nlm.nih.gov/books/NBK305896/#lipid_intro.toc-abstract)>.
68. Bittner V. Lipoprotein abnormalities related to women's health. *Am J Cardiol* 2002; 90(8A): 77i–84i.
69. Phan BA, Toth PP. Dyslipidemia in women: etiology and management. *Int J Womens Health* 2014; 6: 185–194. Dostupné z DOI: <<http://dx.doi.org/10.2147/IJWH.S38133>>.
70. Cífková R, Krajčoviechová A. Dyslipidemia and cardiovascular disease in women. *Curr Cardiol Rep* 2015; 17(7): 609. Dostupné z DOI: <<http://dx.doi.org/10.1007/s11886-015-0609-5>>.
71. Feingold KR, Grunfeld C. Obesity and Dyslipidemia. In: De Groot LJ, Chrousos G, Dungan K et al (eds). *Endotext* [Internet]. MDText.com: South Dartmouth (MA) 2015. Dostupné z WWW: <<https://www.ncbi.nlm.nih.gov/books/NBK305895/>>.
72. Godsland IF. Effects of postmenopausal hormone replacement therapy on lipid, lipoprotein, and apolipoprotein (a) concentrations: analysis of studies published from 1974–2000. *Fertil Steril* 2001; 75(5): 898–915.
73. Lobo RA. Clinical review 27: Effects of hormonal replacement on lipids and lipoproteins in postmenopausal women. *J Clin Endocrinol Metab* 1991; 73(5): 925–930.
74. Goodman MP. Are all estrogens created equal? A review of oral vs. transdermal therapy. *J Women's Health (Larchmt)* 2012; 21(2): 161–169. Dostupné z DOI: <<http://dx.doi.org/10.1089/jwh.2011.2839>>.
75. Smith GI, Reeds DN, Okunade AL et al. Systemic delivery of estradiol, but not testosterone or progesterone, alters very low density lipoprotein-triglyceride kinetics in postmenopausal women. *J Clin Endocrinol Metab* 2014; 99(7): E1306–E1310. Dostupné z DOI: <<http://dx.doi.org/10.1210/jc.2013-4470>>.
76. [Writing Group for the PEPI Trial]. Effects of estrogen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women. The Postmenopausal Estrogen/Progestin Interventions (PEPI) Trial. *JAMA* 1995; 273(3): 199–208. Erratum in *JAMA* 1995; 274(21):1676.
77. Kim JJ, Choi YM. Dyslipidemia in women with polycystic ovary syndrome. *Obstet Gynecol Sci* 2013; 56(3): 137–142. Dostupné z DOI: <<http://dx.doi.org/10.5468/ogs.2013.56.3.137>>.
78. Wild RA. Dyslipidemia in PCOS. *Steroids* 2012; 77(4): 295–299. Dostupné z DOI: <<http://dx.doi.org/10.1016/j.steroids.2011.12.002>>.